Year: 2024 September Volume: 31 Number: 3



Medical Journal of Süleyman Demirel University

Süleyman Demirel Üniversitesi Tıp Fakültesi Dergisi

ISSN 1300-7416 e-ISSN 2602-2109

Dergi

Süleyman Demirel Üniversitesi Tıp Fakültesi

ÜAK Doçentlik başvurularında yer alan "Ulusal Hakemli Dergi" tanımında yer alan ULAKBİM TR Dizin dergi listesidir.

Aramayı Kaydet
Dergi Önerisi

Bulunan Toplam Sonuç : 1

Başlık A-Z

Filtrelerde Ara..

Yıl (16)

- 2024 1
- 2023 1
- 2022 1
- 2021 1
- 2020 1
- 2019 1

Yayın Dili (1)

- İngilizce 1

Konu (54)



Süleyman Demirel Üniversitesi Tıp Fakültesi Dergisi

ISSN: **1300-7416**
EISSN: **2602-2109**
Editör **Prof. Dr. Evrim ERDEMOĞLU**

2024 2023 2022 2021
Tüm Yıllar

Web Adresi

TÜBİTAK ULAKBİM Ulusal Akademik Ağ ve Bilgi Merkezi Cahit Arf Bilgi Merkezi © 2024 Tüm Hakları Saklıdır Kullanım Şartları Gizlilik Politikası

Toplam Kayıt: 1

JSON

İLETİŞİM
TÜBİTAK ULAKBİM TR Dizin
Yüzüncüyıl, İşçi Blokları Mahallesi
Muhsin Yazıcıoğlu Caddesi No:51/C
06530 Çankaya / ANKARA +90 (312) 298 92 00
trdizin@tubitak.gov.tr

Türkiye Cumhuriyeti SANAYİ VE TEKNOLOJİ BAKANLIĞI

TÜBİTAK ULAKBİM

TÜBİTAK ULAKBİM Ulusal Akademik Ağ ve Bilgi Merkezi Cahit Arf Bilgi Merkezi © 2024 Tüm Hakları Saklıdır

p-ISSN 1300-7416
e-ISSN 2602-2109

Medical Journal of Süleyman Demirel University

Süleyman Demirel Üniversitesi Tıp Fakültesi Dergisi

Med J SDU / SDÜ Tıp Fak Derg

The journal is a peer reviewed academic journal and publishes four issues per year in March, June, September, and December.

This journal is indexed by **EBSCO, Tr-Dizin.**

Owner
On Behalf of Suleyman Demirel University
Prof. Dr. Nilgün ŞENOL
Suleyman Demirel University Dean of Faculty of Medicine

Editor in Chief
Prof. Dr. Evrim ERDEMOĞLU (Süleyman Demirel University, Isparta, TÜRKİYE)

Editors
Assoc. Prof. Dr. Sabriye ERCAN (Süleyman Demirel University, Isparta, TÜRKİYE)
Assoc. Prof. Dr. V. Atahan TOĞAY (Süleyman Demirel University, Isparta, TÜRKİYE)

Sectional Editorial Board
Prof. Dr. Mesut ERBAŞ (Çanakkale Onsekiz Mart University, Çanakkale, TÜRKİYE)
Prof. Dr. Ömer ERŞEN (University of Health Sciences, İstanbul, TÜRKİYE)
Prof. Dr. Nilgün GÜRBÜZ (Süleyman Demirel University, Isparta, TÜRKİYE)
Prof. Dr. Pakize KIRDEMİR (Süleyman Demirel University, Isparta, TÜRKİYE)
Prof. Dr. Selma KORKMAZ (Süleyman Demirel University, Isparta, TÜRKİYE)
Prof. Dr. Mustafa NAZIROĞLU (Süleyman Demirel University, Isparta, TÜRKİYE)
Prof. Dr. Aynur AYTEKİN ÖZDEMİR (İstanbul Medeniyet University, İstanbul, TÜRKİYE)
Prof. Dr. Mekin SEZİK (Süleyman Demirel University, Isparta, TÜRKİYE)
Prof. Dr. Coşkun SİLAN (Çanakkale Onsekiz Mart University, Çanakkale, TÜRKİYE)
Prof. Dr. Ali Ekrem ÜNAL (Ankara University, Ankara, TÜRKİYE)
Prof. Dr. Hasan YASAN (Süleyman Demirel University, Isparta, TÜRKİYE)
Assoc. Prof. Dr. Özlem COŞKUN (Çanakkale Onsekiz Mart University, Çanakkale, TÜRKİYE)
Assoc. Prof. Dr. Şeref Selçuk KILIÇ (Çukurova University, Adana, TÜRKİYE)
Assoc. Prof. Dr. Mehtap SAVRAN (Süleyman Demirel University, Isparta, TÜRKİYE)
Assoc. Prof. Dr. Memnun SEVEN (University of Massachusetts Amherst, Massachusetts, USA)
Asst. Prof. Dr. Oruç Numan GÖKÇE (Çanakkale Onsekiz Mart University, Çanakkale, TÜRKİYE)
Asst. Prof. Dr. Buket GÜNGÖR (Çanakkale Onsekiz Mart University, Çanakkale, TÜRKİYE)
Asst. Prof. Dr. Katherine MOORE (Mayo Clinic, Minnessota, USA)
Dr. Katarzyna JAROSZ (International University of Logistics and Transport, Wrocław, POLAND)
Dr. Edin KABİL (University of Sarajevo, Bosnia, BOSNIA AND HERZEGOVINA)
Dr. Ardalan SHARIAT (Tehran University of Medical Sciences, Tehran, IRAN)
Dr. Ülkü YAYLALI ÜSER (Antalya Training and Research Hospital, Antalya, TÜRKİYE)

Language Editor
Asst. Prof. Dr. Muhammed Burak SELVER (Süleyman Demirel University, Isparta, TÜRKİYE)

Statistics Editor
Asst. Prof. Dr. Adnan KARAİBRAHİMOĞLU (Süleyman Demirel University, Isparta, TÜRKİYE)

Graphic Design
Lecturer Serdağ DAĞLI (Süleyman Demirel University, Isparta, TÜRKİYE)

Editorial Office
Dr. Vildan KAYA (Süleyman Demirel University, Isparta, TÜRKİYE)

Address
SDÜ Tıp Fakültesi Dergisi Sekreterliği Isparta, TÜRKİYE

Press
SDÜ Basımevi / ISPARTA

SDÜ Tıp Fakültesi Dekanlığı / 32260 ISPARTA
Tel: 0 246 2113714 - 2113230 - Fax: 0 246 2371165
sdu.tip.dergi@sdu.edu.tr - http://dergipark.gov.tr/sdutfd

Medical Journal of Süleyman Demirel University Authors Guidelines

About Med J SDU
Medical Journal of Süleyman Demirel University (Med J SDU) is published quarterly in March, June, September and December. Med J SDU is an international, scientific, open access, online/print journal that follows independent, unbiased, and double-blind peer-review principles.

Med J SDU publishes research in the health sciences, including original clinical and experimental studies, reviews of current topics, case reports, editorial comments, and letters to the editor. The language of the journal is **English.**

Med J SDU is indexing in both international (EBSCO) and national (TRDizin) indexes.

There is no charges for publishing or publication process. No copyright price are payable to the authors or other third parties for the articles published in the journal. Med J SDU has adopted the policy of providing open access with the publication. Authors' credentials and e-mail addresses are in no way used for other purposes.

The editorial and publication processes of the journal are shaped in accordance with the guidelines of the International Council of Medical Journal Editors (ICMJE). The journal conforms to the Principles of Transparency and Best Practice in Scholarly Publishing (doaj.org/bestpractice).

Originality, high scientific quality and citation potential are the most important criteria for a manuscript to be accepted for publication. Manuscripts submitted for evaluation should not have been previously presented or already published in an electronic or printed medium. Manuscripts that have been presented in a meeting should be submitted with detailed information on the organization, including the name, date, and location of the organization.

Ethical Principles
An approval of research protocols by the Ethics Committee in accordance with international agreements (World Medical Association Declaration of Helsinki "Ethical Principles for Medical Research Involving Human Subjects," amended in October 2013, www.wma.net) is required for experimental, clinical, and drug studies and for some case reports. Ethics committee approvals or an equivalent official documents must be uploaded into the dergipark system.

• For manuscripts concerning experimental researches on humans, a "Written Informed Consent to Participate and Publish" statement should included in the text.

• Written Informed Consent to Participate Publish statement and Ethics Committee approval details (name of the committee, date and number) should included in the Materials and Methods section and in the end of the article (before references) with separate sub-headings.

• It is the authors responsibility to carefully protect the patients anonymity. For photographs that may reveal the identity of the patients, releases signed by the patient or their legal representative should be enclosed.

• For studies carried out on animals, the measures taken to prevent pain and suffering of the animals should be stated clearly. Ethics Committee approval details (name of the committee, date and number) should included in the Materials and Methods section and in the end of the article (before references) with separate sub-headings.

All submissions are screened by a similarity detection software (iThenticate) and the similarity limitation is **25%.** The Editorial Board of the journal handles all appeal and complaint cases within the scope of Committee on Publication Ethics (COPE) guidelines. In such cases, authors should get in direct contact with the editorial office regarding their appeals and complaints. When needed, an ombudsperson may be assigned to resolve cases that cannot be resolved internally. The Editor in Chief is the final authority in the decision-making process for all appeals and complaints.

When submitting a manuscript to Med J SDU, authors accept to assign the copyright of their manuscript to the journal. If rejected for publication, the copyright of the manuscript is considered to be assign back to the authors. Each submission must be submitted in accordance with the journal template (available for download at: https://dergipark.org.tr/tr/download/journal-file/24521), together with a Copyright Transfer Form (available for download at: https://dergipark.org.tr/tr/download/journal-file/22117). Mandatory docu-ments to be sent can be found at https://dergipark.org.tr/tr/pub/sdutfd.

When using previously published content, including figures, tables, or any other material in both print and electronic formats, authors must obtain permission from the copyright holder. Legal, financial and criminal liabilities in this regard belong to the author(s). Statements or opinions expressed in the manuscripts published in Med J SDU reflect the views of the author(s) and not the opinions of the editors, the editorial board, or the publisher; the editors, the editorial board, and the publisher disclaim any responsibility or liability for such materials. The final responsibility in regard to the published content rests with the authors. Revisions should submit within 15 days in pre-review stage and in 30 days in review stage. Otherwise manuscripts will be rejected.

Manuscript Preparation
Manuscripts can only be submitted through the journal's online manuscript submission and evaluation system, available at https://dergipark.org.tr/tr/pub/sdutfd. Manuscripts submitted via any other medium will not be evaluated. Manuscripts submitted to the journal will first go through a technical evaluation process where the editorial office staff will ensure that the manuscript has been prepared and submitted in accordance with the journal's guidelines. Submissions that do not conform to the journal's guidelines will be returned to the submitting author with technical correction requests. The editor reserves the right to reject manuscripts that do not comply with the above-mentioned requirements. Corrections may be done without changing the main text. The editor reserves the right to reject the articles that do not meet the required conditions.

Authors are required to submit the following:
• *Copyright Transfer Form* (Authors must use journal template)

• *Title Page* (Authors must use journal template, must include title of manuscript, running title, authors' name, title and institution, corresponding author's contact information, ORCID addresses, name of the organization supporting the research and ethical approvals)

• *Main document* (It should be prepared according to the journal template and should include a sectioned abstract; Full text file must also include tables and figures with legends alongside separate table and figure files, must be suitable for double blind review process.).

• *Figures* (If applicable, Jpeg/png format, at least 300 DPI, high resolution, up to 6 figures)

• *Tables* (If applicable, Microsoft Word format, up to 6 table, Each table should not exceed one page)

• *Ethical Approvals* (If applicable)

• *Author Contributions* (according to CRediT, please see *Template*, Authors must use journal template)

• Plagiarism Report
• Language Editing Certificate (Certificate should be obtained

from an institution or corporation providing this service or from a person whose main business is language services such as a sworn translator)

Preparation of the Main Document
The articles should be written with double-spaced in 12 pt, Times New Roman character and at least 2 cm from all edges of each page. The main text should not contain any information about the authors' names and affiliations. Original articles should have a structured abstract (Objective, Material and Methods, Results, Conclusion, Keywords). For case reports and reviews, the structured abstract is not used. Limit the abstract to 300 words. References, tables and citations should not be used in an abstract. Authors must include relevant keywords (3-5) on the line following the end of the abstract. All acronyms and abbreviations used in the manuscript should be defined at first use, both in the abstract and in the main text. The abbreviation should be provided in parentheses following the definition. When a drug, product, hardware, or software program is mentioned within the main text, product information, including the name of the product, the producer of the product, and city and the country of the company, should be provided in parentheses.

All references, tables, and figures should be referred to within the main text, and they should be numbered consecutively. The symbols used must be in nomenclature standards.

Original Research Articles should be maximum of 4000 words and include subheadings below;
• Title
• Abstract
• Keywords
• Introduction
• Material and Methods
• Results
• Discussion
• Conclusions
• Statements (Please See: Template)
• References
• Figure and Table Legends (if necessary)

Case Reports should be maximum of 2000 words and include subheadings below;
• Title
• Abstract
• Keywords
• Introduction
• Case Presentation
• Discussion and Conclusion
• Statements (Please See: Template)
• References
• Figure and Table Legends (if necessary)

Reviews should be maximum of 6000 words and include subheadings below;
• Title
• Abstract
• Keywords
• Main text
• Conclusion
• Statements (Please See: Template)
• References
• Figure and Table Legends (if necessary)

Letter to Editor should be maximum of 1000 words and include subheadings below;
• Title
• Abstract
• Keywords
• Letter to Editor
• Statements (Please See: Template)
• References
• Figure and Table Legends (if necessary)

Preparation of the Figures and Tables
• Up to 6 Figures, graphics, or photographs can be submitted. Separate files (in JPEG format) should be submitted through the submission system and should not be embedded in a Word document. The minimum resolution of each submitted figure should be 300 DPI. Figures should be in high resolution so that the printed version can be read easily.

• Figures with legends should also be listed at main document for review.

• Information or illustrations must not permit identification of patients, and written informed consent for publication must be sought for any photograph.

• Up to 6 tables can be submitted. Each table should not exceed one page. Tables should be listed at main document for review.

• All tables and figures should be numbered consecutively.

References
All references should be numbered consecutively in Vancouver style. If more than two consecutive resources are used, only the first and last source numbers should be specified (for example; 2-6). Journal titles should be abbreviated in accordance with the journal abbreviations in Index Medicus. When there are 3 or fewer authors, all authors should be listed. If there are 4 or more authors, the first 3 authors should be listed followed by "et al." The reference styles for different types of publications are presented in the following examples:

For journals;
Neville K, Bromberg A, Bromberg S, el al. The third epidemic multidrug resistant tuberculosis. Chest 1994;1(4):45-8. 555

For books;
Sweetman SC. Martindale the Complete Drug Reference. 34th ed. London: Pharmaceutical Press; 2005.

For book section;
Collins P. Embryology and development, Neonatal anatomy and growth. In: Williams PL, Bannister LH, Berry MM, Collins P, Dyson M, Dussek JE, Ferguson MWJ. Gray's Anatomy (38th Ed) London, Churchill Livingstone, 1995; 91-342.

For website;
Gaudin S. How moon landing changed technology history [Internet]. Computerworld UK. 2009 [cited 15 June 2014]. Available from: http://www.computerworlduk.com/in-depth/it-business/2387/how-moon-landing-changed-technology-history/

For conference proceeding;
Brom A. Robotics and Mechatronics. Symposium on Robotics, Mechatronics and Animatronics in the Creative and Entertainment Industries and Arts. SSAISB 2005 Convention. University of Hertfordshire, Hatfield, UK; 2005.

For Thesis;
Ercan S. Venöz yetmezlikli hastalarda kalf kası egzersizlerinin venöz fonksiyona ve kas gücüne etkisi. Suleyman Demirel University Faculty of Medicine Sports Medicine Department Thesis. Isparta: Suleyman Demirel University. 2016.

Retraction or Reject
Manuscript Retraction: Authors may withdraw their manuscript from the journal with a written declaration with detailed reason only before DOI assigned. If a DOI has been assigned to the article, it can no longer be withdrawn by the authors.

Manuscript Reject: Editors have the authority to reject articles at any stage of the article process.

After Acceptance
If the manuscript is accepted, the acceptance letter and the last version of manuscript are sent to corresponding author through submission system. The journal board is authorized to decide in which volume of the accepted article will be printed. Authors may publish their articles on their personal or corporate websites with the appropriate cite and library rules after publication.

Contents

Research Articles

The Relationship Between Hematological Inflammatory Markers and Postoperative Hypocalcemia in Patients with Primary Hyperparathyroidism
Emre GEZER, Ömer ZEKEY, Büşra YAPRAK BAYRAK, Alev SELEK, Berrin ÇETİNARSLAN, Zeynep CANTÜRK, Damla KÖKSALAN, Mehmet SÖZEN 205

Evaluation of Hepatitis Serology Screening Frequency and Viral Reactivation in Patients Followed with Biological Therapy or Cytotoxic Chemotherapy
Arif KILÇAR, Atalay DOĞRU 213

Fluvoxamine Administration Attenuates Lipopolysaccharide-Induced Pancreatic Damage
Şenay TOPSAKAL, Özlem ÖZMEN 221

An Essential Problem in Patients with Hereditary Angioedema: Irritable Bowel Syndrome
Mehmet KILINÇ, Fatih ÇÖLKESEN, Filiz SADİ AYKAN, Recep EVCEN, Eray YILDIZ, Tuğba ÖNALAN, Mehmet Emin GEREK, Şevket ARSLAN 229

Evaluation of Patients Diagnosed with Spondylodiscitis
Onur ÜNAL, Melahat YILMAZ, Gül Ruhsar YILMAZ, Füsun Zeynep AKCAM, Onur KAYA, Esra Nurlu TEMEL 235

Psychiatric Emergencies in Perinatal Women: A Retrospective Analysis
Kumru ŞENYAŞAR METERELLİYOZ, Alevhan İNAN ÜNLÜ, Pınar ÇETINAY AYDIN 243

Exploring Internalized Stigma, Self-Esteem, and Symptom Severity in Depression: A Comparative Study of Active and Remitted Phases
Pınar ÜNAL AYDIN, Emin Oryal TAŞKIN 253

Perceived Stress in Medical Education: Relationship with Empathy and Stigmatisation
Gözde BACIK YAMAN, Berna NAR, Gökçe İŞCAN 261

Case Reports

Catamenial Hemoptysis: A Case Report and Literature Review
Hasan Emre YILDIRIM, Hasan Ekrem CAMAŞ, Süleyman Emre AKIN, Rasih YAZKAN 271

Reviews

Effects of School-Based Interventions Implemented by Nurses for Children Aged 3-6 Years: A Systematic Review of Experimental Evidence
Gökçe ALGÜL, Ebru KILIÇARSLAN 277

ORIGINAL RESEARCH

Med J SDU / SDÜ Tıp Fak Derg ▸ 2024:31(3):205-211 doi: 10.17343/sdutfd.1413088

The Relationship Between Hematological Inflammatory Markers and Postoperative Hypocalcemia in Patients with Primary Hyperparathyroidism

Emre GEZER[1], Ömer ZEKEY[2], Büşra YAPRAK BAYRAK[3], Alev SELEK[1], Berrin ÇETİNARSLAN[1], Zeynep CANTÜRK[1], Damla KÖKSALAN[1], Mehmet SÖZEN[1]

[1] Kocaeli University, Faculty of Medicine, Department of Endocrinology and Metabolism, Kocaeli, Türkiye
[2] Kocaeli University, Faculty of Medicine, Department of Internal Medicine, Kocaeli, Türkiye
[3] Kocaeli University, Faculty of Medicine, Department of Pathology, Kocaeli, Türkiye

Cite this article as: Gezer E, Zekey Ö, Yaprak Bayrak B, Selek A, Çetinarslan B, Cantürk Z, Köksalan D, Sözen M. The Relationship Between Hematological Inflammatory Markers and Postoperative Hypocalcemia in Patients with Primary Hyperparathyroidism. Med J SDU 2024;31(3):205-211.

Abstract

Objective

Primary hyperparathyroidism (PHPT) is a common endocrine disease that is characterized by hypercalcemia and commonly associated with parathyroid adenoma (PTA). Hypocalcemia is a common postoperative complication in patients with PHPT. The neutrophil-to-lymphocyte ratio (NLR) and platelet-to-lymphocyte ratio (PLR) are inexpensive hematological inflammatory markers. We aimed to investigate the potential predictive risk factors, including the hemogram-derived inflammatory markers for early postoperative hypocalcemia in patients with PHPT.

Material and Method

Patients diagnosed with PHPT, underwent parathyroidectomy and histopathologically shown to be caused by a single PTA were included.

Results

NLR was significantly correlated with parathormone (PTH), while PLR was related considerably with only NLR. A significant positive correlation was shown between gland weight, volume, calcium (Ca), and PTH levels. A significant correlation of postoperative hypocalcemia with age, preoperative Ca, PTH, and NLR was also demonstrated.

Conclusion

We found that NLR was significantly higher in patients with PHPT who developed postoperative hypocalcemia; however, our regression analysis did not find elevated NLR as a significant predictive risk factor for postoperative hypocalcemia. To the best of our knowledge, this is the first study investigating the relationship between hemogram-derived inflammatory markers and clinical parameters, such as the development of postoperative hypocalcemia and preoperative nephrolithiasis, in patients with PHPT.

Keywords: Hyperparathyroidism, hypocalcemia, neutrophils, lymphocytes, platelets

Correspondence: Ö.Z. / dr.zekey@gmail.com
Received: 01.01.2024 • Accepted: 03.07.2024
ORCID IDs of the Authors: E.G: 0000-0002-5340-6106; Ö.Z: 0009-0004-4142-3605; B.Y.B: 0000-0002-0537-3127; A.S: 0000-0002-0646-8697; B.Ç: 0000-0002-8041-8161; Z.C: 0000-0001-7114-2565; D.K: 0000-0002-6349-2976; M.S: 0000-0002-8428-1115

Introduction

Primary hyperparathyroidism (PHPT) is an endocrine disease that is presented with hypercalcemia and is commonly associated with a parathyroid adenoma (PTA) (1). Parathyroidectomy has been described as the gold standard therapy for PHPT to normalize serum calcium (Ca) and parathyroid hormone (PTH) levels (2). Hypocalcemia is a common postoperative complication with a 2.3-42% incidence in patients with PHPT (3,4). Even if postoperative hypocalcemia is most commonly temporary, the outcome of this early postoperative complication might be life-threatening; therefore, hospitalization for several days after parathyroidectomy has been recommended by many experienced physicians (5,6). Multiple factors, such as age, PTH, alkaline phosphatase, preoperative serum calcium, osteocalcin, and vitamin D levels, were found to correlate with postoperative hypocalcemia in patients with secondary hyperparathyroidism. At the same time, there was no correlation between laboratory or other clinical factors and postoperative hypocalcemia in patients with primary hyperparathyroidism (4,7–9).

The neutrophil-to-lymphocyte ratio (NLR) and platelet-to-lymphocyte ratio (PLR) are inexpensive hematological inflammatory markers, and these markers are currently in use to indicate the severity of systemic inflammation or infectious complications (10,11). A relationship between PHPT and systemic inflammation has been described by several studies, as a possible consequence of elevated serum calcium levels, which might trigger systemic inflammation (12,13). Furthermore, PTH alone has been shown to stimulate interleukin-6 production and high PTH levels have been reported in septic patients (14,15). In line with these data, NLR has been found significantly related to various parameters, such as serum PTH, Ca, and P levels, white blood cell count, and parathyroid adenoma size (16–18). In this study, we aimed to investigate the potential predictive risk factors, including the hemogram-derived inflammatory markers for early postoperative hypocalcemia in patients with PHPT.

Material and Method

Patients aged >18 years and who were diagnosed with PHPT, underwent parathyroidectomy and histopathologically shown to be caused by a single PTA at our institute between 2010 and 2021 were included in our retrospective study. Demographics, preoperative albumin-corrected serum Ca, phosphorus (P), PTH, 25-hydroxy vitamin D (25OH-D), 24-hour urine calcium (24hr UCa), neutrophil, platelet, and lymphocyte counts, gonadal status of patients, lowest postoperative corrected calcium, the presence of nephrolithiasis demonstrated by any imaging technique and weight of the surgically removed parathyroid gland were recorded. Additionally, lengths of 3 dimensions of PTA given by our pathology department were recorded to calculate the volume of PTA. The volume of a parathyroid adenoma was calculated using the formula for an ellipsoid (Volume = [4/3] π r1r2r3, where r1, r2, r3 are the lengths of the semiaxes in centimeters). Corrected calcium was derived using the following calculation: serum calcium (mg/dL) + 0.8 x (4 - serum albumin [g/dL]). PLR was calculated as the ratio of platelet count to lymphocyte count and NLR as the ratio of neutrophil count to lymphocyte count. Postoperative hypocalcemia was described as a serum calcium level of less than 8.5 mg/dL on the first biochemical test following parathyroidectomy.

The IBM SPSS for Windows version 20.0 (IBM Corp., Armonk, NY, USA) was used to perform all statistical analyses. The normality assumption was assessed by using the Kolmogorov-Smirnov and Shapiro-Wilk's tests. Continuous variables were given as either mean standard deviation or median (interquartile range) depending on the distribution pattern. The Mann-Whitney U test was used to perform the comparisons between groups and the Spearman's correlation analysis was used to determine the correlations between continuous variables. The receiver operating characteristics (ROC) analysis was used to determine the area under the curve (AUC) and cut-off values. A p-value <0.05 was considered statistically significant.

Results

A total of 368 patients were included in our study. The basic clinical and demographic features of this population are presented in Table 1. The mean age of the patients was 53.1 ± 12.9, with a female-to-male ratio of approximately 5 to 1. The Spearman's correlation analysis demonstrated that NLR was significantly correlated with PTH and PLR ($p = 0.001$, $r = 0.173$; $p < 0.001$, $r = 0.517$, respectively), while PLR was significantly related to only NLR. A significant positive correlation between gland weight and volume, and Ca and PTH levels was shown (weight and Ca, $p = 0.001$, $r = 0.187$; weight and PTH, $p < 0.001$, $r = 0.370$; volume and Ca, $p < 0.001$, $r = 0.198$; volume and PTH, $p < 0.001$, $r = 0.324$), while there was a significant but very weak negative correlation between gland volume and weight, and P levels (weight and P, $p = 0.044$, $r = -0.119$; volume and P, $p = 0.009$, $r = -0.141$). A very strong correlation between gland weight and volume ($p < 0.001$, $r = 0.831$) was also demonstrated.

Table 1 Demographics and clinical characteristics of patients (n=368)

Characteristic	n (%)
Age (years), median (min-max)	54 (18-84)
Gender	
male	60 (16.3)
female	308 (83.7)
Gonadal Status	
menopausal	158 (52.5)
eugonadal	143 (47.5)
Preoperative Nephrolithiasis	
yes	47 (28.3)
no	119 (71.7)
Postoperative Hypocalcemia	
yes	81 (22.1)
no	286 (77.9)
median (25^{th}-75^{th} percentile)	
Ca (mg/dL)	10.7 (10.3-11.2)
P (mg/dL)	2.6 (2.3-3.0)
PTH (pg/mL)	194 (135-305)
Postoperative PTH (pg/mL)	29 (13-74)
25OH-D (ng/mL)	17 (11-25)
NLR	1.95 (1.52-2.69)
PLR	117.45 (90.93-148.02)
Macroscopic Characteristics of Adenoma	
weight (gr)	1.10 (0.50-2.36)
volume (cm^3)	5.86 (2.51-16.03)

Ca: calcium NLR: neutrophil lymphocyte ratio; P: phosphorus; PLR: platelet lymphocyte ratio; PTH: parathyroid hormone; 25OH-D: 25-hydroxy vitamin D

A significant relationship between postoperative hypocalcemia and age, preoperative Ca, PTH, and NLR was demonstrated ($p = 0.001$, $p = 0.014$, $p = 0.003$, and $p = 0.031$, respectively) by the Mann-Whitney U test (Table 2). NLR values of patients who developed postoperative hypocalcemia were significantly higher than patients without. ROC curve analysis was performed to estimate preoperative predictive levels of Ca, PTH, age, and NLR for postoperative hypocalcemia (Figure 1). The threshold Ca level that would distinguish between patients who developed postoperative hypocalcemia and those who did not was 10.3 with an AUC of 0.589 ($p = 0.02$), 351 for PTH with an AUC of 0.614 ($p < 0.01$), 51 for age with an AUC of 0.623 ($p < 0.01$), and 2.95 for NLR with an AUC of 0.579 ($p = 0.03$). Moreover, there was also a significant association between the preoperative nephrolithiasis and preoperative PLR, P, Ca x P, and 24 hr UCa levels ($p = 0.004$, $p = 0.005$, $p = 0.012$, and $p = 0.039$, respectively).

The independent risk factors for hypocalcemia were investigated by using a multivariate linear regression analysis (Table 3). After adjusting for the potential

contributors to postoperative hypocalcemia, i.e., age, parathyroid gland weight, preoperative Ca, PTH, PRL, and 25OH-D levels, the significance of the correlation between NLR and the postoperative hypocalcemia disappeared (p = 0.332). In this multivariate regression analysis, younger age, lower preoperative Ca, and higher PTH levels were shown as the only predictive risk factors for postoperative hypocalcemia (p = 0.022, p = 0.029, and p = 0.013, respectively).

Table 2 The comparison between the clinical characteristics and the early postoperative hypocalcemia in patients with primary hyperparathyroidism

	Postoperative Hypocalcemia		
	Yes¶	No¶	p*
Ca	10.7 (10.1-11.0)	10.8 (10.4-11.2)	**0.014**
P	2.7 (2.2-3.2)	2.6 (2.3-2.9)	0.422
Ca x P	27.8 (24.1-31.0)	27.8 (24.0-31.9)	0.844
PTH	227 (148-987)	187 (133-284)	**0.003**
NLR	2.0 (1.7-3.1)	1.9 (1.5-2.6)	**0.031**
PLR	127.6 (94.3-178.9)	115.7 (88.7-144.2)	0.051
24 hr UCa	392 (201-446)	312 (225-445)	0.764
Age	49.0 (36.5-59.0)	56.0 (45.0-63.0)	**0.001**
Gland weight	1.5 (0.5-2.5)	1.0 (0.5-2.2)	0.660
Gland volume	6.0 (2.5-16.2)	5.9 (2.4-16.5)	0.656

Ca, calcium; NLR, neutrophil lymphocyte ratio; P, phosphorus; PTH, parathyroid hormone; PLR, platelet lymphocyte ratio; 24 hr UCa, 24-hour urinary calcium Bold font indicates statistical significance. αData are expressed as mean ± standard deviation ¶Data are expressed as median (25th-75th percentile) *Evaluated by Mann-Whitney U-test

Table 3 Multiple logistic regression analyses of the adjusted factors, correlation with the early postoperative hypocalcemia in patients with primary hyperparathyroidism.

	OR	95% CI for OR	p
Age	0.963	0.993 - 0.994	0.022
Ca	0.558	0.331 – 0.941	0.029
PTH	1.001	1.000 - 1.002	0.013
NLR	1.232	0.808 – 1.878	0.332
PLR	1.006	0.997 – 1.015	0.190
25OH-D	1.000	0.964 – 1.037	0.995
Gland weight	1.000	1.000 – 1.000	0.329

Ca, calcium; CI, confidence interval; NLR, neutrophil lymphocyte ratio; OR, odds ratio; P, phosphorus; PTH, parathyroid hormone; PLR, platelet lymphocyte ratio; 25OH-D, 25-hydroxy vitamin D Bold font indicates statistical significance




Figure 1:
Receiver operating characteristic curve analysis of preoperative
(a) Ca, (b) PTH, (c) age, and (d) NLR levels for predicting postoperative hypocalcemia.

Discussion

In our study, NLR levels were significantly higher in patients with PHPT who developed postoperative hypocalcemia compared to those who had normal postoperative Ca levels. On the other hand, after adjusting for other confounding factors, the results showed elevated NLR was not significantly a predictive risk factor for postoperative hypocalcemia. To date, this is the first study that investigated the correlation between hemogram-derived inflammatory markers and postoperative hypocalcemia in patients with PHPT. We also showed that younger age, lower serum preoperative Ca levels, and higher PTH levels were predictive risk factors for the development of postoperative hypocalcemia in patients with PHPT. A significant, but very weak correlation between NLR and PTH levels was shown in our analysis. Therefore, it can be stated that PTH may play a role as a confounding factor that causes elevated NLRs in patients with postoperative hypocalcemia compared to those without.

The predictive risk factors for postoperative hypocalcemia which were found in our study are newly discovered findings for the literature. Mittendorf et al. have reported that there was no significant association between any of the laboratory values or other clinical factors, and postoperative hypocalcemia in 162 patients with PHPT (4). In another study with similar sample size, the authors found no significant relationship between preoperative PTH, Ca, P, magnesium, albumin, blood urea nitrogen, thyroid stimulating hormone and vitamin D levels, and postoperative early and permanent hypocalcemia in patients with PHPT (9).

A recent study conducted by Liu et al. supports and contradicts some findings of this present study. In line with our findings, the authors found that the patients who were younger than 45 years had higher Ca decreases after parathyroidectomy than the older patients (19). Moreover, it was reported that the weight of the removed parathyroid gland predicted the amount of Ca drop postoperatively in patients with PHPT. The lesions heavier than 2g had a more significant Ca decrease following parathyroidectomy. In contrast with this finding, our results showed that the removed gland weight was not a predictive risk factor for postoperative hypocalcemia.

In our correlation analysis, we demonstrated a significant but very weak correlation between NLR and preoperative serum PTH levels. Our finding is consistent with the results that were presented by a study with a similar sample size (16). Toraman et al. have reported that NLR showed a significant and very weak correlation with serum PTH, vitamin D, and Ca levels in patients with elevated PTH. Similarly, Zeren et al. demonstrated a significant moderate correlation

between NLR and preoperative serum PTH levels in only 32 patients with PHPT (17). In line with this report, a significant medium correlation between NLR and preoperative serum PTH levels was shown by another study which included 37 patients with PHTP (18). The relatively small sample sizes in these two studies might be the main factors that augmented the correlation levels between these two parameters.

Conclusion

In conclusion, we found that NLR was significantly higher in patients with PHPT who developed postoperative hypocalcemia; however, our regression analysis did not find elevated NLR as a significant predictive risk factor for postoperative hypocalcemia. Moreover, the patients with nephrolithiasis had significantly higher PLR than those without any renal stones. To the best of our knowledge, this is the first study investigating the correlation between hemogram-derived inflammatory markers and clinical parameters, such as the development of postoperative hypocalcemia and preoperative nephrolithiasis, in patients with PHPT. Secondly, in this presented study, younger age, higher PTH, and lower preoperative serum Ca levels were shown as the predictive risk factors for postoperative hypocalcemia in patients with PHPT. We believe that our findings are of paramount importance to the literature on the management of primary hyperparathyroidism.

Conflict of Interest Statement
The authors declare no conflicts of interest.

Ethical Statement
Ethics approval was obtained from the Non-Invasive Clinical Research Ethics Committee of Kocaeli University (Date:25.10.2021, No: GOKAEK 2021/18.19 - 2021/305). The study was conducted in accordance with the principles set forth in the Declaration of Helsinki.

Consent to Participate and Publish
Written informed consent to participate and publish was obtained from all individual participants included in the study.

Funding
No funding was received in support of this research.

Availability of Data and Materials
Data is available on request from the authors.

Authors Contributions
EG: Conceptualization; Formal analysis; Investigation; Validation; Visualization; Writing-original draft.

ÖZ: Data curation; Investigation; Methodology.

BYB: Data curation; Formal analysis; Investigation; Methodology.

AS: Supervision; Writing-review & editing.

BÇ: Supervision; Writing-review & editing.

ZC: Supervision; Writing-review & editing.

DK: Conceptualization; Investigation; Writing-original draft.

MS: Conceptualization; Investigation; Writing-original draft.

References

1. Williams BA, Trites JRB, Taylor SM, Bullock MJ, Hart RD. Surgical management of primary hyperparathyroidism in Canada. J Otolaryngol - Head Neck Surg 2014;43.
2. Muñoz-Torres M, García-Martín A. Primary hyperparathyroidism. Med Clin (Barc) 2018;150(6):226–32.
3. Singh Ospina NM, Rodriguez-Gutierrez R, Maraka S, Espinosa de Ycaza AE, Jasim S, Castaneda-Guarderas A, et al. Outcomes of parathyroidectomy in patients with primary hyperparathyroidism: A systematic review and meta-analysis. World J Surg 2016;40(10):2359–77.
4. Mittendorf EA, Merlino JI, McHenry CR. Post-parathyroidectomy hypocalcemia: Incidence, risk factors, and management. Am Surg 2004;70(2):114–9.
5. Crea N, Pata G, Casella C, Cappelli C, Salerni B. Predictive factors for postoperative severe hypocalcaemia after parathyroidectomy for primary hyperparathyroidism. Am Surg 2012;78(3):352–8.
6. Vasher M, Goodman A, Politz D, Norman J. Postoperative calcium requirements in 6,000 patients undergoing outpatient parathyroidectomy: Easily avoiding symptomatic hypocalcemia. J Am Coll Surg 2010;211(1):49–54.
7. Phimphilai M, Inya S, Manosroi W. A predictive risk score to diagnose hypocalcemia after parathyroidectomy in patients with secondary hyperparathyroidism: A 22-year retrospective cohort study. Sci Rep 2022;12(1).
8. Yang PS, Liu CL, Liu TP, Chen HH, Wu CJ, Cheng SP. Parathyroidectomy decreases neutrophil-to-lymphocyte and platelet-to-lymphocyte ratios. J Surg Res 2018;224:169–75.
9. Kaya C, Tam AA, Dirikoç A, Kılıçyazgan A, Kılıç M, Türkölmez Ş, et al. Hypocalcemia development in patients operated for primary hyperparathyroidism: Can it be predicted preoperatively? Arch Endocrinol Metab 2016;60(5):465–71.
10. Zahorec R. Neutrophil-to-lymphocyte ratio, past, present, and future perspectives. Bratislava Med J 2021;122(7):474–88.
11. Gasparyan AY, Ayvazyan L, Mukanova U, Yessirkepov M, Kitas GD. The platelet-to-lymphocyte ratio as an inflammatory marker in rheumatic diseases. Ann Lab Med 2019;39(4):345–57.
12. Lam HB, Yang PS, Chien MN, Lee JJ, Chao LF, Cheng SP. Association between neutrophil-to-lymphocyte ratio and parathyroid hormone in patients with primary hyperparathyroidism. Arch Med Sci 2019;15(4):880–6.
13. Cheng SP, Liu CL, Liu TP, Hsu YC, Lee JJ. Association between parathyroid hormone levels and inflammatory markers among US adults. Mediators Inflamm 2014;2014.

14. Lind L, Carlstedt F, Rastad J, Stiernström H, Stridsberg M, Ljunggren Ö, et al. Hypocalcemia and parathyroid hormone secretion in critically ill patients. Crit Care Med 2000;28(1):93–9.
15. Verheyen N, Meinitzer A, Grübler MR, Ablasser K, Kolesnik E, Fahrleitner-Pammer A, et al. Low-grade inflammation and tryptophan-kynurenine pathway activation are associated with adverse cardiac remodeling in primary hyperparathyroidism: The EPATH trial. Clin Chem Lab Med 2017;55(7):1034–42.
16. Toraman A, Aras F, Hekimsoy Z, Kursat S. Is there a relationship between parathyroid hormone and neutrophil-lymphocyte or platelet lymphocyte ratio? Acta Endocrinol (Copenh) 2019;15(1):96–101.
17. Zeren S, Yaylak F, Ozbay I, Bayhan Z. Relationship between the neutrophil to lymphocyte ratio and parathyroid adenoma size in patients with primary hyperparathyroidism. Int Surg 2015;100(7–8):1185–9.
18. Alakuş H, Göksu M. Does Parathyroidectomy affect the neutrophil/lymphocyte ratio, a systemic inflammatory marker? Cureus 2021;13(3).
19. Liu C, Tang L, Goel P, Chambers T, Kokot N, Sinha U, et al. A Practical mathematic method to predict and manage hypocalcemia after parathyroidectomy and thyroidectomy. Ann Otol Rhinol Laryngol 2020;129(1):70–7.