Cerebral Palsy and Executive Functions: Inhibitory Control and Cognitive Flexibility Skills

Begüm Çapa Tayyare[1] | Evrim Gerçek[2] | Erbil Dursun[3] | Nur Akçin[4]

https://doi.org/10.1002/pits.22957

1. Department of Occupational Therapy, Vocational School of Health Services, Kocaeli University, Kocaeli, Turkey
2. Department of Speech and Language Therapy, School of Health Sciences, Maltepe University, İstanbul, Turkey
3. Department of Physical Medicine and Rehabilitation, Medical School, Kocaeli University, Kocaeli, Turkey
4. Department of Special Education, Faculty of Education, Marmara University, İstanbul, Turkey
Correspondence Begüm ÇapaTayyare Email: begum_capa@yahoo.com and begum.capatayyare@kocaeli.edu.tr

Keywords: Inhibitory Control, Cognitive Flexibility, Executive Functions, Cerebral Palsy, Children.

Abstract

Background and Aims: Executive functions (EFs) are an umbrella term that includes various cognitive abilities (such as inhibition, planning, goal-setting, monitoring and shifting). There is common agreement that there are three main EFs: inhibitory control, cognitive flexibility, and working memory. Further studies have reported that compared to typically developing peers, children with cerebral palsy (CP) show deficits in EF skills. This study aims to evaluate children with CP's inhibitory control and cognitive flexibility skills.

Methods and Procedures: Forty children with CP between the ages of 7-13 were included in the present study. EF scores of the CP group were compared with scores of typically developing peers matched one-to-one by age and sex. The EF skills were evaluated with Wisconsin Card Sorting Test (WCST), Stroop Color-Word Test (SCWT) TBAG Form, and Executive Function Behavior Rating Scale Parent Form (BRIEF-P). It was determined that there was a significant difference in the WCST total trials, total errors, categories completed, and percent conceptual level responses scores ($p<.05$); SCWT-TBAG scores ($p<.05$) and BRIEF-P all scores ($p\leq.01$) of children with CP.

Conclusions and Implications: The results of this study suggest that children with CP are disadvantaged with EF skills in terms of inhibitory control and cognitive flexibility.

1. Introduction

Cerebral palsy (CP) is a group of disorders that result from non-progressive damage to the developing fetal or infant brain, leading to movement and posture impairments and causing activity limitations (Rosenbaum et al., 2007). Although brain damage is non-progressive, the clinical manifestations may change as a result of age, growth, and central nervous system development (Weierink, Vermeulen, & Boyd, 2013). According to The Surveillance of Cerebral Palsy in Europe, CP is separated into spastic, dyskinetic, ataxic subtypes. It has been observed that approximately 70-80% of individuals with CP exhibit clinical features consistent with the spastic subtype (Krigger, 2006). CP is typically characterized as a motor development disorder, but it encompasses a wide range of impairments beyond movement disorders. Individuals with CP often experience challenges related to sensory perception, cognitive, communicative, and behavioral problems (Fluss & Lidzba, 2020; Bax, Flodmark, & Tydeman, 2007). Disability and functional limitations are often determined by comorbidities such as epilepsy, scoliosis, auditory and visual deficits, mental disability, and behavioral challenges (Shevell, Dagenais &Oskoui, 2013). These non-neuromotor comorbidities can be the primary reason for the burden of care and healthcare usage for individuals with CP and their families (Shevell,2009).

CP has long been considered from a medical perspective, almost special priority is given to motor pathology (Cabezas &Carriedo, 2020). Recently the studies are increasingly highlighting cognitive disorders in children with CP. Recent reviews have reported that children with spastic CP frequently experience challenges with attention and executive function (EF) skills (Stadskleiv, 2020; Stadskleiv, Jahnsen, Anderson, & Tetzchner, 2018; Laporta-Hoyos, et al. 2019; Fluss & Lidzba, 2020). EF skills have been defined as an umbrella term that includes high-level cognitive processes involved in goal-directed, flexible, and adaptive behavior and the top-down regulation of cognition and behavior that are especially actuated in difficult and complex situations (Zelazo, 2020; Roebers, 2017; Goldstein, Naglieri, Princiotta, & Otero, 2014; Freire & Osorio, 2020; Diamond, 2013; Bodimeade, Whittingham, Lloyd & Boydl, 2013; Anderson, 2001). EFs enable individuals to manage and change their actions at their sole discretion considering the consequences of each behavior (e.g. “I can't watch TV more because I need to study for next week's exam”) (Pereira et al., 2018; Botvinick, 2001). And play important roles in purposeful action, emotion regulation, social participation, functional independence, quality of life and success in school (Zelazo, 2020; Voyer, Nadeau & Tessier, 2018; Sakash, et al., 2018; Mousavi, Akbarfahimi, Moein & Vahedi, 2022; Federica & Mellone, 2022; Dawson & Guare, 2014; Bottcher, 2010). EFs, which have many different definitions or features, are mainly examined under three main headings: working memory, inhibitory control and cognitive flexibility (Diamond, 2013).

Working memory is defined as an interface between perception, short-term memory, long-term memory, and target-specific actions (Diamond, 2013). It plays a vital role in oral communication, listening and reading

comprehension, written expression, and correct long-term learning (Meltzer, 2014). Working memory begins to develop in infancy and continues to develop throughout in mid-late childhood (Anderson et al., 2010). Inhibitory control is defined as the central component of EFs (Barkley, 1997). Generally, it focuses on the ability to actively inhibit or delay a dominant response to achieve a goal (Morasch & Bell, 2011). It involves controlling attention, behavior, thoughts, or emotions and doing what is more appropriate or necessary to override a strong inner predisposition or an attractive external stimulus (e.g. suppressing an impulsive utterance) (Zelazo, 2020; Diamond, 2013). Inhibitory control begins to develop from the end of the first year of age and continues to develop rapidly across the toddler period and into the preschool years (Morasch & Bell, 2011; Diamond, 2006; Anderson et al., 2010). Cognitive flexibility is closely related to creativity and changing setup. It represents the ability to adapt to changing conditions. The ability to change plans in the face of obstacles, setbacks, new information, or faults (Dawson & Guare, 2014). Cognitive flexibility develops rapidly in the preschool period in parallel with the development of neural networks of the prefrontal cortex, and this development continuously increases well into adolescence. (Buttelman & Karbach, 2017).

Studies show that children with spastic CP have deficits in inhibitory control (Li et al., 2014; Christ et al., 2003; Cabezas & Carriedo, 2020; Caillies, Hody, & Calmus, 2012; Bottcher, Flachs & Uldall, 2010), conitive flexibility (Crichton et al., 2020; Bodimeade, Whittingham, Lloyd & Boydl, 2013), planning (Stadskleiv et al., 2014; Bodimeade, Whittingham, Lloyd & Boydl, 2013), working memory (Van Rooijen, Verhoeven & Steenbergen, 2016; Løhaugen et al., 2014; Di Lieto et al., 2017), and information processing (Laporta-Hoyos, et al. 2019; Bottcher, Flachs & Uldall, 2010). On the other hand, children with dyskinetic CP tend to have mild EF deficits (Fluss & Lidzba, 2020) and normal comprehension and visual perception (Stadskleiv, 2020). Some of these studies have evaluated the EF skills in children with CP using clinical measures and compared them to typically developing peers (Laporta-Hoyos, et al. 2019; Freire & Osorio, 2020; Crichton et al., 2020; Bodimeade, Whittingham, Lloyd & Boydl, 2013). Some of them have evaluated EF skills using behavior rating scales (in daily life skills) (Sørensen et al., 2016; Muriel et al., 2015; Laporta-Hoyos, et al. 2022). Only a few studies have used both clinical measures and behavioral ratings of EF together (Whittingham, Bodimeade, Lloyd, & Boyd, 2014; Mousavi, et al., 2022; Bottcher, Flachs & Uldall, 2010). Additionally, several studies have investigated the effectiveness of various interventions including computer-based cognitive training and physical exercise in improving the EF performance of children with CP (Piovesana, et al. 2017; Pereira, 2019; Lakes et al., 2019; García-Galant, et al. 2020; Di Lieto et al., 2021). While the studies have reported improvements in EF skills following these interventions, further research is needed to determine the most effective interventions and to better understand the underlying mechanisms of EF deficits in CP.

Poor inhibitory control and mental flexibility skills can be impacts individuals with CP in all aspects of daily life. For example Roostaei et al. (2021) highlight the challenges that people with CP face when engaging in dual-tasking activities. Dual-task refers to the ability to perform two or more tasks simultaneously, which is a common occurrence in our daily lives (Okur et al., 2022; McIsaac et al., 2015). The tasks could be either motor or cognitive tasks (Huang & Mercer, 2001). Inhibitory control enables individuals to suppress automatic or prepotent responses, which is important for successful dual-task performance (Verbruggen &Logan, 2008). The sensory-motor disorders as a result of early brain damage may often expose children with CP to dual tasks (Reilly, et al., 2008). The systematic review and meta-analysis of observational studies show that verbal fluency tasks are more likely to lead to a decline in walking speed compared to discrimination/decision-making tasks and mental tracking tasks (Such as maintaining a straight posture while reading) (Roostaei et al., 2021). The study concluded that the effects of dual-task conditions were more pronounced in children with CP, and that walking and balance performance deteriorated under these conditions. This makes it necessary to look at CP from a multi-faceted perspective.

The International Classification of Functioning, Disability, and Health (ICF) (Word Health Organization, 2007) recommends considering personal and environmental factors as well as body structure and functions for activity and participation. In the traditional CP rehabilitation model, poor attention is paid to the cognitive characteristics of individuals (Cabezas &Carriedo, 2020). Extending patient outcomes beyond sensory-motor disorders to include activity and participation is a valuable and patient-centered approach to CP treatment (Eslinger, 2022), as it is recommended by ICF. Evaluation of EFs in CP will allow the individual's rehabilitation plan to be formulated in a way that will affect all aspects of life as recommended in the ICF. Therefore, clinicians working with CP should consistently ask about non-motor disabilities for appropriate assessment, rehabilitation, and guidance (Fluss & Lidzba, 2020).

In summary individuals with CP are at risk for EF deficits, which can have a significant impact on their daily functioning. Early identification and intervention can help to improve EF skills and mitigate the impact of CP on cognitive functioning. In the narrative review of cognitive and academic profiles in children with CP, Fluss & Lidzba (2020) suggest that in high-risk infants or those diagnosed with early CP, regular general evaluation from early childhood to late childhood, when cognitive impairment may occur. They also suggest that elements of attention and executive deficit should be sought with the neuropsychological assessment tools to offer early guidance to parents and teachers who are often unprepared and confused by children with CP. It is also important to note that EF deficits in CP are not a homogeneous group. Different types of CP and different levels of impairment can result in varying degrees and patterns of EF deficits. Therefore, personalized assessment and intervention strategies are needed to address the specific needs of each individual with CP. Further research is needed to better understand the underlying mechanisms of EF deficits in CP and to identify the most effective interventions for this population (Sakash, et al., 2018; Mousavi, Akbarfahimi, Moein & Vahedi, 2022; Cabezas & Carriedo, 2020).

In this context, the main aim of our study is to determine whether the neuropsychological assessment (Inhibitory control and mental flexibility) and daily life skills of EFs of children with spastic CP are different from the reference group. Additionally, the aim is to establish whether the correlation findings between these two measurements provide clinical insight for evaluating EF in children with CP.

2. Method

2.1 Participants

A total of 130 children and adolescents with CP who had received a rehabilitation program at the Physical Medicine and Rehabilitation clinic of Kocaeli University Medical Faculty were evaluated. Children diagnosed with spastic CP between the age of 7-13, who volunteered with the permission of their parents, were included. Children who could not read, did not know primary colors, had severe dystonia or epileptic seizures, and had mental retardations or other neurodevelopmental disorders were excluded. The EFs scores of the children with spastic CP (CP group) were compared with those of reference group. The CP group was matched one-to-one by age and sex with their typically developing peers in the normative data. The reference group was established using the EF normative data scores of Turkish children. Table 1 presents the demographic characteristics of the CP group.

Table 1. Demographic data

	Age (Mean±SD)	**Gender (n/%)**	**CP Type (n/%)**	**GMFCS* (n/%)**
CP Group	9,4±1,9	18 (%45) F 22 (%55) M	7 hemiplegic(%17,5) 23 diplegic (%57,5) 10 quadriplegic (%25)	I (n=9 %22,5) II (n=11 %27,5) III (n=8 %20) IV (n=12 %30)
Total		n=40	n=40	n=40

* Gross Motor Function Classification System: I: Walks without Limitations, II: Walks with Limitations, III: Walks Using a Hand-Held Mobility Device, IV: Self-Mobility with Limitations; May Use Powered Mobility

To evaluate the EF skills, Wisconsin Card Sorting Test (WCST), Stroop Color-Word Test (SCWT) TBAG Form, and Executive Function Behavior Rating Scale Parent Form (BRIEF-P) were used. The study protocol was approved by the Kocaeli University Faculty of Medicine Ethics Committee with the project number GOKAEK 2017/155 and dated 07/06/2017.

2.2 Materials

The study involves EFs evaluating measurement tools in the BİLNOT-Children neuropsychological test battery (Karakaş & Doğutepe-Dinçer, 2011), which was standardized for a sample of children in Turkey to provide standardized neuropsychological tests. The tests were conducted by a researcher certified in administering and scoring the tests.

Wisconsin Card Sorting Test (WCST)

WCST consist of four stimulus cards containing three stimulus parameters (color, shape and number). Participants are required to sort the numbered answer cards according to different principles and change their approach during test administration (Grant & Berg, 1948). Responses were classified as: correct errors, persistent responses and non-perseverative errors. The WCST is primarily used to assess perseveration and abstract thinking (Grant & Berg, 1948). According to Greve et al. (2005), WCST is a high-level measurement tool that measures EFs. The WCST Turkish Form is associated with property such as attention,

feature determination, perseveration, working memory, executive functions, conceptualization, and abstract thinking (Karakaş & Doğutepe-Dinçer, 2011). WCST contains four stimulus cards and one hundred twenty-eight response cards (Figure 1). The cases are required to find the correct category by the given feedback that the response is true or false (without mentioning the category name) while taking WCST (Erdoğan-Bakar et al., 2011).

Within the scope of this study, total trials (W1), total errors (W2), perseverative responses (W5), perseverative errors (W6), percent perseverative errors (W8), trials to complete first category (W9), which measure inhibition, and the number of correct responses (W4), conceptual level responses (W10), and percent conceptual level responses (W11) scores that measure set-up maintaining were calculated. The scores calculated from the responses to WCST are as in Figure 2.




Figure 1. WCST (Wisconsin Card Sorting Test)

WCST Scores	
W1	Total trials
W2	Total errors
W3	Total number of correct responses
W4	Number of categories completed
W5	Perseverative responses
W6	Perseverative errors
W7	Total number of non- perseverative errors
W8	Per cent perseverative errors
W9	Trials to complete the first category
W10	Conceptual level responses
W11	Percent conceptual level responses

Figure 2. Scores of WCST

Stroop Color-Word Test (SCWT) TBAG Form

The Stroop Color-Word Test TBAG Form, which is a neuropsychological measurement tool, is basically an EF test (Karakaş, 2011). It measures the ability to change the perceptual setup in line with changing demands and, under a "disruptive effect", suppress a habitual behavioral pattern and perform an unusual behavior, as well as focused attention (Scarpina & Tagini, 2017; Anderson, 2001). The SCWT-TBAG form was formed by combining the original Stroop Color-Word Test and the Victoria Form (Karakaş ve Doğutepe-Dinçer, 2011).




Figure 3. Stroop Color-Word Test (TBAG Form)

Sections	Stimulus	Scope of the stimulus card	Task
1.Section	1.Card	Color names printed in black ink	Read color name
2.Section	2.Card	Color names printed in an inconsistent color ink	Read color name
3.Section	3.Card	Colored dots	Say dot color
4.Section	4.Card	Neutral words printed in color ink	Say ink color of word
5.Section	2.Card	Colors printed in an inconsistent color ink	Say ink color of word

Figure 4. Stroop Color-Word Test (TBAG Form)

In the SCWT-TBAG Form, subjects are required to read/say five task as fast as possible (Figure 4.). Firstly, participants are required to read the names of colors printed in black ink (1. card in figure 3). Secondly, participants are required to read the names of colors printed in an inconsistent color ink (2. card in figure 3). The third table represents colored dots. Thirdly participants are required to say the color of circles (shape-color: SC). The fourth table represents inconsistent words which printed in an inconsistent color ink (For example, the word "weak" in red ink). In the fourth table, participants are required to say the ink colors of the words. The fifth table is the same as the second table (the word “yellow” is printed in blue ink). Fifthly, participants are required to say the color of the ink instead of reading the word (named color-word: NCW).

The SCWT has various scoring systems. In this study, we used named color-word fifth section scores (time, number of errors and corrections) to measure the inhibition. In addition, the difference 3 (Dif. 3) score was also used in the study to exclude the speaking speed factor. Dif.3 score was calculated by subtracting the time score of named shape-color from the time score of named color-word (Difference between NCW and SC).

Behavior Rating Inventory for Executive Functions (BRIEF)

BRIEF (Gioia et al., 2000) can be applied to children between the ages of 5-18. The validity of the scale in Turkish culture was carried out by Erdoğan-Bakar and friends (2011). There are two forms of BRIEF, teacher and parent. Both forms consist of 86 items and 8 subscales that include EF behaviors. Because the assessment is made in different environments, 18 items of the teacher and parent forms are different from each other. Three indicator scores are calculated from these subscales. The first is the "Behavior Regulation Index (Inhibit, Emotional Control, and Shift)", the second is the "Metacognition Index (Initiate, Working Memory, Plan/Organize, Organization of Materials, Monitor)", and the third is the "Global Executive Composite" obtained from the sum of these two indicator scores (Figure 5). The parent form of BRIEF (BRIEF-P) has been used for this study. Parents indicate if the child shows difficulties concerning the behavior which given the test item with "1 (never)", "2 (sometimes)", or "3 (often)". High scores on BRIEF suggests poor EF skills (Erdoğan Bakar et al.,2011; Gioia et al, 2000) .




Figure 5. The test scores of executive functioning

2.3 Statistical Evaluation

Statistical evaluation was made with IBM SPSS 20.0 (IBM Corp., Armonk, NY, USA) package program. The normal distribution of the scores of the sample was evaluated by the Kolmogorov-Smirnov test. Since the samples did not show a normal distribution, the difference between the groups was evaluated with the Mann-Whitney U test. Correlation analyzes were evaluated by Spearman. For the testing of two-way hypotheses, $p < 0.05$ was considered sufficient for statistical significance.

3. Results

The scores on WCST total trials ($p = 0.00$), total errors ($p = 0.04$), categories completed ($p = 0.00$) and percent conceptual level responses ($p = 0.03$) scores were statistically significant. There was no significant difference in the scores of perseverative responses, perseverative errors, percent perseverative errors, trials to complete first category and conceptual level responses ($p > 0.05$) (Table 2.).

Table 2. WCST scores

	CP Group (Mean±SD)	Reference Group (Mean±SD)	p
Total trials	123,5 ±12,28	113,82 ±14,28	**.000**
Total Errors	55,6 ±22,42	45,55 ±14,17	**.040**
Categories Completed	3±1,94	4,21±0,94	**.001**
Perseverative Responses	37,9±32,68	31,18±10,84	.729
Perseverative Errors	32,35±24,09	26,98±8,93	.893
Percent Perseverative Errors	25,59±18,65	22,21±5,91	.350
Trials to Complete 1st Category	31,4±34	15,72±4,56	.996
Conceptual Level Responses	50,95±21,75	55,01±5,94	.544
Percent conceptual Level Responses	42,51±20,7	51,54±12,7	**.038**

The SCWT-TBAG and BRIEF-P test scores of the children with CP significantly differed ($p<0.05$) from the reference group (Table 3., Table 4.).

Table 3. SCWT-TBAG scores

	CP Group (Mean±SD)	Reference Group (Mean±SD)	p
S5Time (S5T) (sec.)	73,97±37,17	40,63±6,86	**.000**
S5Error (S5E)	1,9±2,25	0,73±0,12	**.020**
S5Correction (S5C)	3,25±2,32	2,04±0,45	**.004**
Difference 3 (S5T-S3T)	38,6±24,46	22,54±4,32	**.002**

Table 4. BRIEF-P scores

Indicator	CP Group (Mean±SD)	Reference Group (Mean±SD)	p
Behavioral Reg.Index (BRI)	48,72±7,98	43,6±2,44	**.010**
Metacognition Index (MI)	83,67±17,87	70,97±2,63	**.000**
Global Executive Composit (GEC)	132,4±24,03	114,61±3,9	**.000**

There was a positive correlation between SCWT-TBAG scores with BRIEF-P BRI ($p < 0.05$), MI ($p < 0.01$), and GEC ($p < 0.01$) and WCST W2 ($p < 0.01$) scores (Table 5.).

Table 5. Correlation between SCWT-TBAG scores with BRIEF-P and WCST scores

		SCWT-TBAG	
		S5Time r */p	Dif. 3 r */p
BRIEF-P	BRI	,346/**.029**	,381/**.015**
	MI	,428/**.006**	,406/**.009**
	GEC	,434/**.005**	,428/**.006**
WCST	Total trials	,318/**.045**	,308/.053
	Total Errors	,464/**.003**	,405/**.009**
	Categories Completed	-,366/**.020**	-,305/.056
	Perseverative Responses	,236/.138	,248/.122
	Perseverative Errors	,267/.095	,266/.095
	Percent Perseverative Errors	,266/.097	,257/.109
	Trials to Complete 1st Category	,261/.103	,224/.164
	Conceptual Level Responses	-,401/**.010**	-,335/**.035**
	Percent Conceptual Level Responses	-,448/**.004**	-,389/**.013**

*Spearman's correlation

There was a negative correlation between SCWT-TBAG scores with WCST Conceptual level responses (S5T $p=0.01$; Difference3 $p < 0.05$) and percent conceptual level responses (S5T $p < 0.01$; Difference3 $p < 0.05$) scores. It was observed that there was no correlation ($p > 0.05$) between perseverative responses, perseverative errors, and percent perseverative errors, trials to complete 1st category scores. While there was a positive correlation between SCWT-TBAG S5T score with WCST total errors ($p < 0.05$) score, a negative correlation between SCWT-TBAG S5T score with WCST categories completed ($p < 0.05$) score and there was no correlation between SCWT-TBAG Difference 3 score and WCST categories completed score (Table 5.).

Table 6. Correlation between BRIEF-P and WCST scores

		BRIEF-P		
		BRI r */p	MI r */p	GEC r */p
WCST	Total trials	,339/**.002**	,470/**.000**	,448/**.000**
	Total Errors	,231/**.039**	,417/**.000**	,301/**.007**
	Categories Completed	-,331/**.003**	-,396/**.000**	-,409/**.000**
	Perseverative Responses	,202/.073	,254/**.023**	,201/.074
	Perseverative Errors	,203/.071	,268/**.016**	,214/.057
	Percent Perseverative Errors	,168/.136	,180/.111	,254/.172
	Trials to Complete 1st Category	,067/.557	,224/.046	,195/.083
	Conceptual Level Responses	-,079/.485	-,131/.247	-,048/.673
	Percent Conceptual Level Responses	-,210/.062	-,415/**.000**	-,282/**.011**

*Spearman's correlation

Table 6 presents correlations between the WCST and BRIEF-P scores. The WCST total trials and total errors scores correlated with the BRIEF-P BRI ($p < 0.05$), MI ($p < 0.01$) and GEC ($p < 0.01$) scores positively. The WCST categories completed score and BRIEF-P BRI ($p < 0.05$), MI ($p < 0.01$) and GEC ($p < 0.01$) scores

negatively correlated. The WCST perseverative responses and perseverative errors correlated with BRIEF-P MI score positively ($p < 0.05$). The WCST percent conceptual level responses correlated with BRIEF-P MI ($p < 0.01$) and GEC ($p < 0.05$) scores negatively.

4. Discussion and Conclusion

The results of this study suggest that children with spastic CP have problems with EF skills in terms of inhibitory control and cognitive flexibility. The significant difference in “completed categories” and “percent conceptual level responses” scores of WCST show that children with spastic CP have problems in maintaining the setting. Nadeau, Routhier & Tessier (2008) reported similar results in the study that compared the WCST performance of children with spastic CP aged 9-12 with typical development peers (control group, n=50). Unlike our findings, they reported "trials to complete 1st category” score of children with spastic CP was higher than the control group.

Inhibitory control enables individuals to regulate and adapt their behavior and conform to established social norms (Federica & Mellone, 2022). The lower performance of children with spastic CP in SCWT-TBAG and WCST “total trials, total errors, categories completed” scores reflected the difficulties with inhibitory control. Freire & Osorio (2020) have also reported differences in inhibitory control between preschool children with CP and typically developing peers.

As far as the findings obtained from BRIEF-P, in which EF skills are evaluated by parents through daily life activities, children with spastic CP experience more difficulties than their peers with typical development in terms of behavior regulation (Inhibit, emotional control and shift); metacognition indicators (Initiate, working memory, plan/organize, organization of materials, monitor) and global executive composite. These findings are similar to past researches showing that children with spastic CP have EF difficulties (Whittingham et al., 2014; Sorensen et al., 2016; Sakash, et al., 2018; Nadeau, Routhier & Tessier, 2008). Dissimilar to the other studies, we examined the correlation between BRIEF-P scores with WCST and SCWT-TBAG scores. The results of correlations reflected that fewer problems with inhibition and impulsivity are associated with lesser executive problems in daily life.

According to the constructivist models that have emerged with the rapid developments in neuroscience in recent years, maturation is a matter of self-structuring mental development through the interaction between innate structures and environmental configuration (Arsaliodu & Pacual-Leone, 2016). Researchers describe the immature nervous system as a highly moldable system that hosts innate biological constraints and also through processing the active, ongoing environmental input and environmental input of the child (Bottcher, 2010). Early brain lesions are not the only reason for the difficulties that children with CP experience in their EF skills. Motor dysfunctions resulting from this damage limit children's learning environment and social environment during their developmental period. Some deficiencies will inevitably arise due to limited learning environments during the construction of mental processes when no intervention is made for children with CP for EF skills, which are thought to mature with age.

A systematic review of interventions for preventing and treating children with CP (Novak & et al. 2020) has reflected that cognitive interventions are useful in a rehabilitation program and the tendency is to include those cognitive interventions to the rehabilitation programs. EFs play an important role in behavior regulation, social skills, and performance of daily living activities, as well as in the regulation of other cognitive skills (Garcia-Galant et al.,2020). Zelazo & et. al (2016) reported that researches clearly show that EF skills can be developed with relatively short interventions, such as increasing the difficulty levels of the tasks. Therefore, children with CP should be supported with various intervention programs regarding EF skills. Considering all these and the EF disorders in our study, for an effective rehabilitation program, it might be beneficial to evaluate all cognitive-linguistic, academic, social, and emotional characteristics of children with CP to determine the situations that create activity limitations (Defined in ICF): “Difficulties an individual may have in executing activities.”] and make treatment arrangements accordingly.

Children with CP are unique, just like every child. Besides the location, effect, and size of the lesions, which compose different combinations of activity limitations and participation in daily living skills of children with CP, the personality and context of the children also form another combination. The critical point here should be the use of neuropsychological tests in a way that allows for recognition of the individual's neuropsychological profile for the design of intervention programs that have functional goals for activities and participation.

5. Limitations

In this study, the inhibitory control and cognitive flexibility skills of children with spastic CP were evaluated with neuropsychological tests. Functions such as learning, visio-spatial perception, decision-making, and working memory that can be measured by different neuropsychological tests were not evaluated.

The other limitation of this study was the lack of the IQ scores of the children with CP. Fluency reading ability was assumed to be the indicator of standard intelligence for this study. According to previous research, standard intelligence tests have limitations in measuring IQ scores of children with CP because some subtests depend on motor skills (Coceski et al., 2021; Sherwell et al., 2014; Yin Foo, Guppy & Johnston, 2013). More studies are needed to measure the intelligence of children with motor impairment.

Acknowledgements

This study is a part of the thesis titled "Executive Function Levels of Children with Cerebral Palsy and Effectiveness of Neuro-Feedback Applications" and the project was supported by the Kocaeli University Scientific Research Projects Unit. Project number: 2018/066. We would like to acknowledge the financial support provided by The Scientific Research Projects Unit of Kocaeli University, which made this research possible.

References

Aisen, M. L., Kerkovich, D., Mast, J., Mulroy, S., Wren, T. A., Kay, R. M., & Rethlefsen, S. A. (2011). Cerebral palsy: clinical care and neurological rehabilitation. *The Lancet Neurology*, *10*(9), 844-852.

Anderson, V. (2001). Assessing executive functions in children: biological, psychological and developmental considerations. *Pediatric Rehabilitation, 4*(3), 119–136.

Anderson, V., Spencer-Smith M., Coleman, L., Anderson, P., Williams, J., Greenham, M.,….Jacobs, R. (2010). Children's executive functions: Are they poorer very early brain insult. *Neuropsychologia, 48*(7), 2041-2050.

Arsaliodu M ve Pacual-Leone J. (2016). Constructivist developmental theory is needed in developmental neuroscience. *Science of Learning, 16016*. doi:10.1038/npjscilearn.2016.16

Barkley, R.A. (1997). Behavioral inhibition sustained attention, and Executive Functions: Constructing a unifying theory of ADHD. Psychological Bulletin, 121 (1), 65-9.

Bax, M.C.O., Flodmark, O. & Tydeman, C. (2007). From syndrome toward disease. *Developmental Medicine & Child Neurology, 109,* 39-41.

Bodimeade, H.L., Whittingham, K., Lloyd, O. & Boyd, R.N. (2013). Executive functioning in children with unilateral cerebral palsy: protocol for a cross-sectional study. *BMJ Open accessible medical research, 3*.

Bottcher, L. (2010) Children with spastic cerebral palsy, Their cognitive functioning, and social participation: A review. *Child Neuropsychology*, *16*, 209–228. doi:10.1080/09297040903559630

Bottcher, L., Flachs, E.M., & Uldall, P. (2010). Attentional and executive impairments in children with spastic cerebral palsy. *Developmental Medicine&Child Neurology*, *52*, 42-47. doi.org/10.1080/09297040903559630

Botvinick, M.M., Braver, T.S., Barch, D.M., Carter, C.S. & Cohen, J.D. (2001). Conflict monitoring and cognitive control. *Psychological Review, 108*(3), 624–52. doi.org/10.1037/0033-295X.108.3.624

Buttelman, F. & Karbach, J. (2017). Development and Plasticity of Cognitive Flexibility in Early and Middle Childhood. *Frontiers Psychology, 36.*

Cabezas, M. & Carriedo, N. (2020). Inhibitory control and temporal perception in cerebral palsy. *Child Neuropsychology*, *26:3,* 362-387, DOI: 10.1080/09297049.2019.1656712

Caillies, S., Hody, A., & Calmus, A. (2012). Theory of mind and irony comprehension in children with cerebral palsy. *Research in developmental disabilities*, *33*(5), 1380-1388.

Christ, S. E., White, D., Brunstrom, J. E., & Abrams, R. A. (2003). Inhibitory control following perinatal brain injury. *Neuropsychology, 17(1),* 171

Chung, H.J., Weyandt, L.L. & Swentosky, A. (2014). The Physiology of Executive Functioning. Goldstein, S. & Naglieri, J.A. (Ed.) *Handbook of Executive Functioning*. New York: Springer Science+Business Media

Coceski, M., Hocking, D.R., Reid, S.M., Abu-Rayya, H.M., Reddihough, D.S., Wrennall, J. & Stargatt, R. (2021). Assessing IQ in adolescents with mild to moderate cerebral palsy using the WISC-V. *The Clinical Neuropsychologist*, 1-20

Crichton, A., Ditchfield, M., Gwinn, S., Wallen, M., Thorley, M., Bracken, J.,…Hoare, B. (2020). Brain magnetic resonance imaging is a predictor of bimanual performance and executive function in children with unilateral cerebral palsy. *Developmental Medicine & Child Neurology*, *62*(5), 615-624.

Dawson, P., & Guare, R. (2014). Interventions to Promote Executive Development in Children and Adolescents. Goldstein, S. & Naglieri, J.A. (Ed.) *Handbook of Executive Functioning*. New York: Springer Science+Business Media.

Diamond, A. (2013). Executive Functions. *Annual review of psychology*, *64,* 135–168.

Diamond, A. (2006). The early development of executive functions. Bialystok E. & Craik, F.I.M. (Ed.), *Lifespan Cognition: Mechanisms of Change*. New York, NY: Oxford University Press

Di Lieto, M. C., Pecini, C., Brovedani, P., Sgandurra, G., Dell'Omo, M., Chilosi, A. M., ... & Cioni, G. (2021). Adaptive Working Memory Training Can Improve Executive Functioning and Visuo-Spatial Skills in Children With Pre-term Spastic Diplegia. *Frontiers in Neurology*, *11*, 601148.

Di Lieto, M.C., Brovedani, P., Pecini, C. Chilosi, A.M., Belmonti, V., Fabbro, F.,...Cioni, G. (2017). Spastic diplegia in preterm-born children: executive function impairment and neuroanatomical correlates. *Research in Developmental Disabilities, 61*, 116-126

Edgin, J.O., Inder, T.E., Anderson, P.J., Hood, K.M., Clark, C.A.C., & Woodward, L.J. (2008). Executive functioning in preschool children born very preterm: Relationship with early white matter pathology. *Journal of the International Neuropsychological Society*, *14*, 90–101.

Erdoğan Bakar, E., Işık Taner, Y. Soysal, A.Ş., Karakaş, S., Turgay, A. (2011). Behavioral rating inventory and laboratory tests measure different aspects of executive functioning in children. *Bull Clin Psychopharmacol;21(4):*302-16.

Eslinger, P. (2022). Investigating language, social cognition, and executive function in cerebral palsy outcomes. *Developmental medicine and child neurology*, 10.1111/dmcn.15483. Advance online publication. https://doi.org/10.1111/dmcn.15483

Federica, F. & Mellone, M. (2022). Executive Function Development: The Mediating Role of Social and Environmental Context. López-Ramón, M.F. (Ed.). *Handbook of Research on Neurocognitive Development of Executive Functions and Implications for Intervention.* Hershey: IGI Global.

Fluss, J., & Lidzba, K. (2020). Cognitive and academic profiles in children with cerebral palsy: a narrative review. *Annals of Physical and Rehabilitation Medicine, 63,* 447–456

Freire, T.C. & Osorio, A.A.C. (2020). Executive Functions and Drawing in Young Children with Cerebral Palsy: Comparisons with Typical Development. Child Neuropsychology, doi:10.1080/09297049.2019.1694648

Garcia-Galant, M., Blasco, M., Reid, L., Pannek, K., Leiva, D., Laporta-Hoyos, O., & et al.(2020) Study protocol of a randomized controlled trial of home-based computerized executive function training for children with cerebral palsy. *BMC pediatrics 20:9*

Gioia, G.A., Isquith, P.K., Guy, S.C., & Kenworthy, L. (2000) Test Review Behavior Rating Inventory of Executive Function. *Child Neuropsychology, 6*(3), 235-238.

Goldstein. S., Naglieri, J.A., Princiotta, D., & Otero, T.M. (2014). Introduction: A History of Executive Functioning as a Theoretical and Clinical Construct. Goldstein, S. & Naglieri, J.A. (Ed.) *Handbook of Executive Functioning*. New York: Springer Science+Business Media

Grant, D. A., & Berg, E. A. (1948). Wisconsin Card Sorting Test. *APA PsycTests*.

Greve, K.W., Stickle, T.R., Love, J.M., Bianchini, K.J., & Stanford, M.S. (2005). Latent structure of the Wisconsin Card Sorting Test: a confirmatory factor analytic study. *Archives of Clinical Neuropsychology*, *20*(3), 355-364.

Huang, H. J., & Mercer, V. S. (2001). Dual-task methodology: applications in studies of cognitive and motor performance in adults and children. *Pediatric Physical Therapy*, *13*(3), 133-140.

Karakaş, S., & Karakaş, H.M. (2000). Yönetici işlevlerin ayrıştırılmasında multidisipliner yaklaşım: bilişsel psikolojiden nöroradyolojiye. *Klinik Psikiyatri, 3*, 215-227.

Karakaş, S. (2011). *Stroop Testi TBAG Formu: Araştırma Geliştirme Çalışmaları ve Kullanım Kılavuzu*. Ankara :Ayrıntı Basım Yayım ve Matbaacılık.

Karakaş, S., & Doğutepe-Dinçer, E. (2011). *BİLNOT-Çocuk Nöropsikolojik Testlerin Çocuklar için Araştırma ve Geliştirme Çalışmaları.* Ankara:Nobel Tıp

Kolk, A., & Talvik, T. (2000). Cognitive Outcome of children with early-onset hemiparesis. *Journal of child neurology*, *15*(9), 581-587.

Krigger, K.W. (2006). Cerebral Palsy: An Overview. *American Academy of Family Physicians*,*73*(1),91-100.

Laporta-Hoyos, O., Ballester-Plané, J., Leiva, D., Ribas, T., Miralbell, J., Torroja-Nualart, C., ... & Pueyo, R. (2019). Executive function and general intellectual functioning in dyskinetic cerebral palsy: comparison with spastic cerebral palsy and typically developing controls. *European Journal of Paediatric Neurology*, *23*(4), 546-559.

Lakes, K. D., Sharp, K., Grant-Beuttler, M., Neville, R., Haddad, F., Sunico, R., ... & Radom-Aizik, S. (2019). A six week therapeutic ballet intervention improved gait and inhibitory control in children with cerebral palsy—a pilot study. *Frontiers in public health*, *7*, 137.

Laporta-Hoyos, O., Ballester-Plané, J., Leiva, D., Ribas, T., Miralbell, J., Torroja-Nualart, C., ... & Pueyo, R. (2019). Executive function and general intellectual functioning in dyskinetic cerebral palsy: comparison with spastic cerebral palsy and typically developing controls. *European Journal of Paediatric Neurology*, *23*(4), 546-559.

Laporta-Hoyos, O., Pannek, K., Pagnozzi, A. M., Whittingham, K., Wotherspoon, J., Benfer, K., ... & Boyd, R. N. (2022). Cognitive, academic, executive and psychological functioning in children with spastic motor type cerebral palsy: Influence of extent, location, and laterality of brain lesions. *European Journal of Paediatric Neurology*, *38*, 33-46.

Li, X., Wang, K., Wu, J., Hong, Y., Zhao, J., Feng, X.,…Zhang, X. (2014). The Link between impaired theory of mind and executive function in children with cerebral palsy. *Research in Developmental Disabilities*, *35*(7), 1686-1693.

Løhaugen, G. C., Beneventi, H., Andersen, G. L., Sundberg, C., Østgård, H. F., Bakkan, E., ... & Skranes, J. (2014). Do children with cerebral palsy benefit from computerized working memory training? Study protocol for a randomized controlled trial. *Trials*, *15*(1), 1-9.

McIsaac, T. L., Lamberg, E. M., & Muratori, L. M. (2015). Building a framework for a dual task taxonomy. *BioMed research international*, *2015*.

Meltzer, L. (2014). Teaching Executive Functioning Processes: Promoting Metacognition, Strategy Use, and Effort. Goldstein, S. & Naglieri, J.A. (Ed.) *Handbook of Executive Functioning*. New York: Springer Science+Business Media.

Mok, M. Y., Yeung, M. K., Chan, Y. S., Lam, C. C., & Chan, D. F. (2019). Neural correlates of attentional control and working memory in children with cerebral palsy: A functional MRI study. Research in Developmental Disabilities,

95, 103486.
Morasch, K.C. & Bell, M.A. (2011). The role of inhibitory control in behavioral and physiological expressions of toddler executive function. *Journal of Experimental Child Psychology*, *108,* 593-606
Mousavi, E., Akbarfahimi, N., Moein, S. & Vahedi, M. (2022). A Study of the Relationship between Executive Function and School Function in Children with Cerebral Palsy. *Journal of Occupational Therapy, Schools, & Early Intervention,* https://doi.org/10.1080/19411243.2022.2027840
Muriel, V., Garcia-Molina, A., Aparicio-Lopez, C., Ensenat, A., & Roig-Rovira, T. (2015). Relationship between executive functioning and behaviour in children with cerebral palsy. *Revista de Neurologia*, *61*(8), 337-343.
Nadeau, L., Routhier, M.E. & Tessier, R. (2008). The performance profile on the Wisconsin Card Sorting Test of a group of children with cerebral palsy aged between 9 and 12. *Developmental Neurorehabilitation*, *11*(2), 134–140.
Novak, I., Morgan, C., Fahey, M., Finch-Edmondson, M., Gales, C.,...&Badawi, N. (2020). State of the Evidence Traffic Lights 2019: Systematic Review of Interventions for Preventing and Treating Children with Cerebral Palsy. *Current Neurology and Neuroscience Reports, 20 (3).*
Okur, E. O., Arik, M. I., Okur, I., Gokpinar, H. H., & Gunel, M. K. (2022). Dual-task training effect on gait parameters in children with spastic diplegic cerebral palsy:Preliminary results of a self-controlled study. *Gait&Posture*,*94*,45-50.
Pacini, A.K., Chevignard, M., Lancien, S., Escolanog, S., Laurent-Vannier, A., Agostinii, M., & Meyer, P. (2017). Executive function after severe childhood traumatic brain injury – age – at-injury vulnerability periods: the TGE prospective longitudinal study. *Annals of Physical and Rehabilitation Medicine, 60*(2), 74-82.
Peng, Y., Zheng, Y., Ding, J., & Zhou, J. (2018). Neural correlates of executive function in children with cerebral palsy: A fMRI study. Journal of Intellectual Disability Research, 62(6), 495-503
Pereira, A., Lopes, S., Magalhaes, P., Sampaio, A., Chaleta, E., & Rosario, P. (2018). How Executive Functions are evaluated in Children with cerebral palsy? A Systematic Review. *Frontiers in Psychology. 9*(21).
Piovesana, A.M., Ross, S., Lloyd, O., Whittingham, K., Ziviani, J., Ware, R. S., & Boyd, R. N. (2017). Randomized controlled trial of a web-based multi-modal therapy program for executive functioning in children and adolescents with unilateral cerebral palsy. *Disability and Rehabilitation*, *39*(20), 2021-2028.
Reilly, D. S., Woollacott, M. H., van Donkelaar, P., & Saavedra, S. (2008). The interaction between executive attention and postural control in dual-task conditions: children with cerebral palsy. *Archives of physical medicine and rehabilitation*, *89*(5), 834-842.
Roebers, C.M. (2017). Executive function and metacognition: Towards a unifying framework of cognitive self-regulation. *Developmental Review, 45*, 31-51.
Roostaei, M., Raji, P., Morone, G., Razi, B., & Khademi-Kalantari, K. (2021). The effect of dual-task conditions on gait and balance performance in children with cerebral palsy: A systematic review and meta-analysis of observational studies. *Journal of bodywork and movement therapies*, *26*, 448-462.
Rosenbaum, P., Paneth, N., Leviton, A., Goldstein, M. & Bax, M. (2007). A report: the definition and classification of cerebral palsy April 2006. *Development Medicine - Child Neurology*, *109*, 8-14.
Sakash, A., Broman, A.T., Rathouz, P.J., & Hustad, K.C. (2018). Executive Function in School-aged children with cerebral palsy: relationship with speech and language. *Research in Developmental Disabilities, 78,* 136-144.
Scarpina F. & Tagini S. (2017) The Stroop Color and Word Test. *Frontiers in psychology. 8*, article 557
Sherwell, S., Reid, S. M., Reddihough, D.S., Wrennall, J., Ong, B., & Stargatt, R. (2014). Measuring intellectual ability in children with cerebral palsy: Can we do better? *Research in Developmental Disabilities 35*(10), 2558-2567
Shevell M, Dagenais L, Oskoui M. (2013) The epidemiology of cerebral palsy: new perspectives from a Canadian registry.Semin Pediatr NeurolJun;20(2):60-4. doi: 10.1016/j.spen.2013.06.008. PMID: 23948680.
Shevell, M. I. (2009). Current understandings and challenges in the management of cerebral palsy. Minerva pediatrica, 61(4), 399-413.
Skranes, J., Evensen, K.I., Lohaugen, G.C., Martinussen, M., Kulseng, S., Myhr, G.,…Brubakk, A. (2008). Abnormal cerebral MRI findings and neuroimpairments in very low birth weight (VLBW) adolescents. *European Journal of Paedıatrıc Neurology*, *12*(4), 273–283.
Sorensen, K., Liverod, J.R., Lerdal, B., Vestrheim, I.E., & Skranes, J. (2016). Executive functions in preschool children with cerebral palsy –Assessment and early intervention – A pilot study. *Developmental Neurorehabilitation, 19*(2), 111–116.
Stadskleiv, K. (2020). Cognitive functioning in children with cerebral palsy. *Developmental Medicine & Child Neurology. 62*, 283–289.
Stadskleiv, K., Jahnsen, R., Anderson, G.L. & Tetzchner, S. (2018). Neuropsychological profiles of children with cerebral palsy. *Developmental Neurorehabilitation, 21*, 2, 108-120.
Stadskleiv, K., von Tetzchner, S., Batorowicz, B., van Balkom, H., Dahlgren-Sandberg, A., & Renner, G. (2014). Investigating Executive Functions in Children with Severe Speech and Movement Disorders Using Structured Tasks, *Frontiers in psychology* doi:10.3389/ fpsyg.2014.00992
Van Rooijen, M., Verhoeven, L., & Steenbergen, B. (2016). Working memory and fine motor skills predict early numeracy performance of children with cerebral palsy. *Child Neuropsychology*, *22*(6), 735-747.
Verbruggen, F., & Logan, G. D. (2008). Automatic and controlled response inhibition: associative learning in the go/no-go and stop-signal paradigms. *Journal of Experimental Psychology: General*, *137*(4), 649.
Voyer, A. P., Nadeau, L., & Tessier, R. (2018). Social dominance in children with cerebral palsy during a problem-solving task with peers. *Disability and rehabilitation*, *40*(19), 2288-2292.
Weierink, L., Vermeulen, R.J., & Boyd, R.N., (2013). Brain structure and executive functions in children with cerebral palsy: A systematic review. *Research in Developmental Disabilities*, *34*(5), 1678-1688.

Whittingham, K., Bodimeade, H.L., Lloyd, O. & Boyd. R.N. (2014) Everyday psychological functioning in children with unilateral cerebral palsy: does executive functioning play a role? *Developmental Medıcıne & Child Neurology, 56*(6), 572–579.

World Health Organization. (2007). *International Classification of Functioning, Disability, and Health: Children & Youth Version: ICF-CY*. World Health Organization.

Yin Foo, R., Guppy, M. & Johnston, L.M. (2013). Intelligence assessments for children with cerebral palsy: a systematic review. *Developmental Medıcıne & Child Neurology, 55*(10), 911–918.

Zelazo, P. D. (2020). Executive function and psychopathology: A neurodevelopmental perspective. *Annual review of clinical psychology*, *16*(1), 431-454.

Zelazo, P. D., Blair, C. B., and Willoughby, M. T. (2016). Executive Function: Implications for Education (NCER 2017-2000). Washington, DC: National Center for Education Research, Institute of Education Sciences, U.S. Department of Education.